



ILUVIEN[®]
(fluocinolone acetonide
intraocular implant) 0.19mg



The next steps of your ILUVIEN treatment journey

APPROVED USE

ILUVIEN is a prescription medicine that is an implant injected into the middle of the eye (vitreous) and is used for the treatment of:

- diabetic macular edema (DME) in patients who have been previously treated with a course of corticosteroids and did not have a significant increase in intraocular pressure.

IMPORTANT SAFETY INFORMATION

- **Do not use ILUVIEN if you have or think you might have:**
 - an infection in or around the eyes, including most viral diseases of the cornea and conjunctiva, including active herpes viral infection of the eye, vaccinia, varicella, mycobacterial infections, and fungal diseases.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information. You may also scan the QR code.





1. Evaluation: Are you a candidate?

You may be a candidate for ILUVIEN if you have been diagnosed with diabetic macular edema (DME).

ILUVIEN is used for the treatment of DME in patients who have been previously treated with a course of corticosteroids and did not have a significant increase in intraocular pressure.

Your eye doctor will perform a detailed evaluation to assess your condition and determine if you are a candidate for ILUVIEN.

For more information visit
ILUVIEN.com

2. Insurance Benefit Investigation

Your retina specialist can submit a benefits investigation with your insurance to evaluate what amount your insurance will pay and the cost to you. If you are unable to pay the amount your insurance does not cover, there may be several options to supplement your existing insurance.

IMPORTANT SAFETY INFORMATION (cont)

- **Do not use ILUVIEN if you have or think you might have: (cont)**
 - glaucoma with a cup to disc ratio of greater than 0.8.
 - a hypersensitivity to any of the ingredients in ILUVIEN.
- **Warnings**
 - Injections into the gel-like substance between the lens and retina of the eye (vitreous) are associated with a serious eye infection (endophthalmitis), eye inflammation, increased or decreased eye pressure, glaucoma, and choroidal or retinal detachments. Your eye doctor should monitor you regularly after the injection.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information. You may also scan the QR code.



3. ILUVIEN Injection


A single ILUVIEN implant is administered by a retina specialist during an in-office procedure. Prior to the procedure, your eye is cleaned and numbing solution is applied. The ILUVIEN implant is injected into the eye. As it is being injected, you may feel some pressure in your eye.

After the injection, if your eye becomes red, sensitive to light, painful, or develops a change in vision, you should seek immediate care from your doctor.

4. Follow-up

Your doctor will monitor your eye after the injection for possible side effects, including development of cataracts, eye infection (endophthalmitis), increased eye pressure, glaucoma, and retinal detachments.

ILUVIEN is designed to
release medicine for up to
36 months.



IMPORTANT SAFETY INFORMATION (cont)

• Warnings (cont)

- If your eye becomes red, sensitive to light, painful, or you notice a change in vision, seek immediate care from your doctor.

For more information visit
ILUVIEN.com

Please see additional Important Safety Information throughout and accompanying full Prescribing Information. You may also scan the QR code.



Questions to ask your doctor about ILUVIEN®

ILUVIEN®
(fluocinolone acetonide
intraocular implant) 0.19mg

1. How does diabetes affect diabetic macular edema (DME)?
2. Are there steps I can take to help slow the progression of DME?
3. How does recurring DME affect my vision?
4. What are my options for treating DME and improving my vision?
5. I have had diabetes for ____ year(s). Will this affect DME treatment types available to me?
6. How does ILUVIEN work?
7. Can I still get ILUVIEN even though I have another eye condition or prior eye surgery?
8. What is the procedure for administering ILUVIEN?
9. What type of payment or insurance coverage is available for ILUVIEN?
10. What are the risks and possible side effects associated with ILUVIEN?
11. What is my follow-up plan after an ILUVIEN injection?



.....

IMPORTANT SAFETY INFORMATION (cont)

• Warnings (cont)

- You may experience temporary blurred vision after receiving an injection and should not drive or use machinery until your vision has resolved.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information. You may also scan the QR code.



For more information visit
ILUVIEN.com

Financial Assistance



ILUVIEN® AccessPlus Financial Assistance Services

If your eye doctor prescribes ILUVIEN, AccessPlus Financial Assistance Services may be able to help find options to assist you with your out-of-pocket costs for patients with or without insurance.

Co-Pay Program

The Co-Pay Program provides co-pay assistance for eligible commercially insured patients with out-of-pocket costs that exceed \$25.

The program is only available for patients with commercial or private insurance and that covers ILUVIEN for the FDA-labeled indication. The patient must be a resident of the US and must provide household income and number in household. The Program does not apply to the patient's cost sharing for the injection procedure, diagnostic tests, or other costs associated with the administration of ILUVIEN.

You or your eye doctor can contact AccessPlus by calling 1-844-445-8843.

Patient Assistance Program

The Patient Assistance Program can provide free treatment for eligible uninsured and rendered uninsured patients.

To be eligible, patients must be residents of the US or Puerto Rico, been prescribed

ILUVIEN for an FDA-approved indication, and patient's household income meets financial restrictions.

To apply for the Patient Assistance Program, complete the ILUVIEN Patient Assistance Program application with your eye doctor. If you are determined eligible, you and your eye doctor will receive an approval letter indicating that you may proceed with scheduling your injection appointment.

Independent Financial Assistance Foundations

Independent financial assistance foundations are independent entities that provide financial assistance for patients insured by a government payer. Contact AccessPlus today to learn more. Call 1-844-445-8843

For more information regarding financial assistance, visit iluvien.com/support/financial-assistance/

Please see Important Safety Information throughout and accompanying full Prescribing Information. You may also scan the QR code.



Important Safety Information

ILUVIEN[®]
(fluocinolone acetonide
intraocular implant) 0.19mg

IMPORTANT SAFETY INFORMATION

(cont)

• Warnings (cont)

- Prolonged use of corticosteroids may result in the development of glaucoma and can cause changes in vision. Your eye doctor should regularly monitor your eye pressure. You may develop increased eye pressure and it may need to be managed with eye drops or surgery.
- The use of corticosteroids may cause: posterior subcapsular cataracts; delay corneal wound healing after cataract surgery; corneal and scleral thinning; and bacterial, viral, or fungal infections. Let your doctor know if you have a history of herpes viral infections of the eye as corticosteroids are not recommended in these patients.
- If the back part of the lens of your eye is missing or torn, the ILUVIEN implant may move to the front part of the eye.

• Common side effects when using ILUVIEN for Diabetic Macular Edema

- The most common side effects in the eye were: cataracts, increased eye pressure, specks that float in the field of vision, eye pain, bleeding in the conjunctiva, clouding of the back of the eye's lens capsule, eye irritation, separation of the gel-like substance inside of the eye (vitreous) from the retina, conjunctivitis, corneal swelling, foreign body sensation in eyes, itchy sensation in the eye, eye redness, damage to the optic nerve, eye discomfort, sensitivity to light, yellow or white spots on your retina, presence of white blood cells in the fluid behind the cornea, and eye discharge.
- The most common side effects not related to the eye were: anemia, headache, kidney failure, and pneumonia.

These are not all of the possible side effects of ILUVIEN. Call your doctor or pharmacist for medical advice about side effects. You may report side effects to Alimera Sciences, Inc. at 1-844-445-8843 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information. You may also scan the QR code.





ILUVIEN[®]
(fluocinolone acetonide
intravitreal implant) 0.19mg



**Ask your doctor if ILUVIEN
is right for you**



Visit [ILUVIEN.com](https://www.iluviens.com) to learn more. Use your mobile device to scan the QR code.

Please see Important Safety Information throughout and accompanying full Prescribing Information. You may also scan the QR code.



© Copyright 2025 ANI Pharmaceuticals, Inc.
All rights reserved. ILUVIEN is a registered trademark of Alimera Sciences Inc., an ANI Pharmaceuticals, Inc. Company.
US-IL-2400012 05/2025



ani
Pharmaceuticals, Inc.